

AD-A098 408

FOREIGN TECHNOLOGY DIV WRIGHT-PATTERSON AFB OH  
ADVANCEMENTS IN CHINESE GEOMAGNETISM AND AERONOMY DURING THE LA--ETC  
FEB 81 Z GANSKUN  
FTD-ID(RS)T-1982-80


F/G A/14  
LA--ETC

UNCLASSIFIED

NL

1981  
FEBRUARY

END
DATE
FILMED
5 81
DTC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A098408

DTIC FILE COPY

FTD-ID(RS)T-1952-80

2

## FOREIGN TECHNOLOGY DIVISION




ADVANCEMENTS IN CHINESE GEOMAGNETISM AND AERONOMY  
DURING THE LAST THIRTY YEARS

by

Zhu Gangkun

DTIC  
ELECTE  
MAY 04 1981  
S D E



Approved for public release;  
distribution unlimited.

81 3 17 057

# EDITED TRANSLATION

FTD-ID(RS)T-1952-80

9 February 1981

MICROFICHE NR: FTD-81-C-000106

ADVANCEMENTS IN CHINESE GEOMAGNETISM AND  
AERONOMY DURING THE LAST THIRTY YEARS,

By: Zhu/Gangkun

English pages: 27

Source: Acta Geophysica Sinica, 1979, pp 326-335

Country of origin: China

Translated by: SCITRAN  
F33657-78-D-0619

Requester: FTD/WE

Approved for public release; distribution  
unlimited.

Accession For	
NTIS GPO-I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION  
FOREIGN TECHNOLOGY DIVISION  
WP.AFB, OHIO.

141

ADVANCEMENTS IN CHINESE GEOMAGNETISM AND AERONOMY  
DURING THE LAST THIRTY YEARS

by Zhu Gangkun

(Institute of Geophysics, Academia Sinica)

Abstract

This paper gives a systematic account of some advancements in the fields of geomagnetism and aeronomy in China during the last thirty years. The review covers the following 14 headings: (1) Establishment of geomagnetic observatories; (2) Geomagnetic surveys and charts; (3) Geomagnetic transient variation and geomagnetic storm prediction; (4) Rock magnetism, paleomagnetism, and archeomagnetism; (5) Magnetotelluric prospecting and related problems; (6) Magnetic prospecting and interpretation of data; (7) Research on geomagnetic instruments; (8) Investigations and theoretical research of seismo-magnetic relations; (9) Statistical analyses of solar-terrestrial relations; (10) Structure of upper atmosphere and the determination of ozone layer; (11) Ionospheric measurements and analyses; (12) Structure and Disturbance of magnetosphere; (13) Studies on cosmic ray intensity variations:

(14) Retrospect and prospect.

[Text]

In the thirty years since the founding of the People's Republic of China, China has made great advancements in geomagnetism and aeronomy, yet the rate of progress has been unbalanced. Generally speaking, progress has been relatively slow and there is still a large gap with the internationally advanced nations. At present, all of China is working hard to realize the "four modernizations" and it is necessary to review the advancements in Chinese geomagnetism and aeronomy over the last thirty years and summarize our own positive and negative experiences.

1. Establishment of Geomagnetic Observatories

Before liberation, China only had the Yushan Observatory near Shanghai which was originally built by a French church. The Central Academy of Physics established a geomagnetic observatory in Nanking for a short time and during the war of resistance against Japan they set up an observatory at Yanshan in Guilin which was later moved to Chongqing. After liberation, under the enthusiastic promotion of the Academia Sinica's Institute of Geophysics, eight geomagnetic observatories were established: Peking Observatory (formal recording started in 1957), Lanzhou

Observatory (1959), Yushan Observatory (1877), Canton Observatory (1957), Changchun Observatory, Lhasa Observatory and Urumqi Observatory. Besides these, to meet the need for geomagnetic measurement, there were geomagnetic observatories provisionally established in Tunhuang and Zhiya.

After the large earthquake at Xingtai in 1966, to actively coordinate earthquake prediction, under the strong support of the Academia Sinica and the National Earthquake Bureau, three key geomagnetic observatories were added at Hongshan in Longyao and at Qingguang in Tianjin. In the last several years, following the establishment of earthquake units in each province and city, various types of survey equipment began to be used all over China and the number of key observatories reached to over 141.

Among the geomagnetic observatories mentioned above, for the first eight established and the two observatories set up since the cultural revolution, the layouts of the observatory sites were homogeneous which developed the basis of geomagnetic observatories. Aside from the Urumqi Observatory, all of the other seven observatories issue formal recordings<sup>[1]</sup> and among them the quality of the Yushan and Peking Observatories is excellent. The other observatories are in the process of vigorous improvement.

## 2. Geomagnetic Surveys and Charts

Prior to 1936, geomagnetic surveying was all done by foreigners in China. Intermittently, 1,128 measurements were done<sup>[2]</sup>.

Thirteen years before liberation, China began to carry out its own geomagnetic surveys and 127 surveys were done on the shores of the Yangtze River, on the Xisha Islands and Hainan Island and in the southwest. After liberation, under the guidance of the Geophysics Institute of the Academia Sinica, geomagnetic surveying increased tremendously. This work can be divided into three general stages. The first stage was from 1950 to 1959 during which time 235 points were surveyed; the second stage lasted until the beginning of the 1970's in which the range was larger and there were initial investigations of 1,740 spots; the third stage was significant in its surveying of long term changing points, the number was limited to under 100 and the area of Tibet was emphasized.

The main work of drawing up Chinese geomagnetic charts was done by the Geophysics Institute of the Academia Sinica. Based on the geomagnetic data collected within China, they compiled Chinese geomagnetic charts of 1950, 1960 and 1970. Most important were the D,I,H,Z value lines and the variation rate charts for other years which could provide a reference for national defense and industrial departments. During these three periods of Chinese geomagnetic charts, they naturally took the chart of 1970 as the best because they used data from more surveyed points (1,740), they gathered time variation patterns from seven Chinese and three Japanese geomagnetic observatories and the space distribution of surveyed points was more homologous than in the past.



At the same time, a written summary of the "analysis method of geomagnetic basic magnetic fields" was also completed [3]. From now on, the direction of work should be focused on the application of aerial surveys and satellite materials and we should begin developing research on long term variations and the generalized method.

### 3. Geomagnetic Transient Variations and Geomagnetic Storm Prediction

Before liberation, China had still not developed this area of research. The mass of geomagnetic data collected at the Yushan Observatory was suitable for analytical research of geomagnetic transient variation. For this, the Geophysics Institute of the Academia Sinica began work in the following areas in 1953:

(1) statistical analyses of geomagnetic storms. A geomagnetic storm catalogue of the Yushan Observatory was completed for the years 1908-1956. The classification of geomagnetic storms was done on the basis of the properties of the violent starts and main phases of geomagnetic storms and statistics were made for the properties of each type of geomagnetic storm including the range of violent starts, the strength of geomagnetic storms and their season and the frequency of sunspot cycles. (2) They completed a K measurement index of the Yushan Observatory for the period 1922-1956 and carried out preliminary statistical analyses.

Recently, detailed work has been done on the two above mentioned projects. <sup>[4]</sup> Geomagnetic bulletins are gradually developing in the

geomagnetic observatories and have been providing timely references. Besides these, they have also determined curved magnetic disturbances, magnetic static sun and magnetic disturbance diurnal variation, and geomagnetic tides [5-6].

In the autumn of 1958, "magnetic storm, ionospheric disturbance and other methods of prediction" were first introduced and based on the "geophysical reaction of sun proton explosions", [7] related units of the Academia Sinica began routine predictions from sun activity and magnetic storms. It was practically proven that not only do magnetic storm predictions rely on effective information of sun activity but it is also necessary to pay close attention to the variations of the high altitude magnetosphere. To perfect and quantize magnetic prediction, it is necessary to expand and deepen work [8].

In recent years, from the interstellar shock wave-geomagnetic storm coupling relation which appeared from solar eruptions, statistical analysis was done on the interrelationship of solar eruptions and geomagnetic storms and an attempt was made to explain these theoretically [8a,8b].

#### 4. Rock Magnetism, Paleomagnetism and Archeomagnetism

Chinese determination of rock specimen magnetism was begun early during the period 1951-1955 by each unit who coordinated magnetic prospecting and mainly determined the magnetism of rocks. Based on the size and other variations of rock and mineral magnetism, we could completely integrate the measured magnetic

force abnormalities and the geological structures of the prospected areas. For earlier work, the reader is referred to references [9-12]. At present, this work is concentrated in the areas of geology, oil and metallurgy. Paleomagnetic measurement and research has been involved with the interpretation of the terrestrial pole position of each geological age and their shifts, the reversals of geomagnetic fields and crustal changes as well as the applications of stratigraphy and geotectology. In early work, reference [13] was based on three samples collected from Gansu and they determined the paleoterrestrial pole as being from the Silurian Period. Reference [14] used a non-directional magnetometer to determine the natural magnetic strength of 175 rock samples and calculated their paleoterrestrial pole position. These samples were collected from north, south and southwest China and were classified as being from five geological eras: the Devonian Period, the Carboniferous Period, the Permian Period, the Triassic Period and the Jurassic Period. In recent work, reference [15] used the alternating demagnetization method to carry out magnetic stability tests on Sinian system sandstone directional samples from Anhui's Xiuning District and gave their average magnetic strength direction and paleoterrestrial pole position. In a similar way, reference [16] carried out paleogeomagnetic research on two basalt section samples from the Nanking area whose geological era is being debated. In recent years, attention has been placed on the paleogeomagnetic determining of

rock samples from the Tibetan plateau and the results are in the process of being collated. Reference [17] used the paleogeomagnetic method to measure for the first time the age of rocks transformed by man in Yunnan. They are about 1 million 700 thousand years old which is earlier than both Peking man and Lantian man. China is a nation with a very long history and civilization and using the magnetism of ancient samples (pottery, bricks, tiles, ancient relics etc.) they developed excellent conditions for archeogeomagnetic work. Reference [18] collected 40 ancient brick samples from the Han dynasty, Three Kingdoms Period and the Tang, Song, Jin, Yuan, Ming and Qing dynasties, processed 427 cubic samples and carried out measurement and gradual heating research on the magnetism of these samples. Thus, they initially determined the geomagnetic dip angle in the Peking area over the last 2,000 years and the geomagnetic field strength and their variations. Archeogeomagnetic work similar to that in the Peking area can without a doubt be extended to samples from other ancient cities and it is predicted that results will soon be published.

##### 5. Magnetotelluric Prospecting and Related Problems

Soon after liberation, when China was searching for oil gas structures and volcanic ores, the use of the natural electric field method was stressed. <sup>[19-21]</sup> At the same time, Chinese scholars published a series of articles [references 22-31] on various ellipsoid natural electric fields, data was collated on geoelectric

currents and there was discussion on the problem of faults. Magnetotellurgic prospecting developed from the geoelectric current method began to be looked at seriously in China at the end of the 1960's and during the 1970's. Reference [32] emphasized the area from Wudu in Gansu Province to Yinchuan in the Ningxia Autonomous Region and at the same time the recorded time of geoelectric magnetic field disturbances was a few seconds to a few thousand seconds. From the measured results, they analyzed the northern section of China's north-south earthquake zone, the earth's crust, the electric features of the upper earth mantle and the electric diversity in earthquake and non-earthquake areas. In the last few years, related fields have continued to use the magnetotellurgic depth-sounder method to research the electric structure of certain crusts in Tibet. The results await publication. Besides these, reference [33] used records of bend disturbances and magnetic storm violent starts from the Peking, Changli, Tianjin and Dalian Geomagnetic Observatories and investigated the conductance abnormalities on the west shore of the Bohai Sea. Reference [34] stressed the records and analysis of 1-200 minute transient geomagnetic variations and abnormalities and their relations to earthquakes from six observatories in the eastern part of Gansu Province. Reference [35] gave a preliminary summary of the basic principles of magnetotellurgic prospecting.

## 6. Magnetic Prospecting and Interpretation of Data

Lately, magnetic measurement on the earth's surface and aero-magnetic measurement have been given serious attention by industrial departments. Its superiority was shown in its being able to attain a type of large scale geological concept in a short period of time for a tremendous surface area. China is a nation of many mountains and its mountainous areas occupy two-thirds of its area. After large scale work, the aero-magnetic measurement method has also made new advancements. Early, in 1966, reference [36] discussed several problems in the research of metallic ore area terrestrial magnetism. References [37-41] stressed the determination and explanation of magnetic abnormalities. In the last ten or more years, the development of magnetic prospecting and the interpretation of data has been mainly concentrated in industrial departments. Following the use of earth resources satellite photos and the United States plan to send up a geomagnetic satellite (Magsat) around September of this year, large area magnetic abnormalities will be discovered and explained and there will no doubt be extensive development.

## 7. Research on Geomagnetic Instruments

After liberation, there was great development in Chinese geomagnetic instruments. There were geological instruments such as the WCWI-63 model non-directional magnetometer, the CHD 5-72 model terrestrial nuclear turning magnetometer, the CCM-1 model

CCK-1

magnetic open gate magnetometer<sup>1</sup> and various other resistance surveying instruments. In a radio factory in Anhui Province's Liuan District, there was also recently produced the QFXC-75 weight nuclear turning magnetometer. The Geophysics Institute of the Academia Sinica has developed the CB<sub>2</sub> triple quantity geomagnetic recorder and the CJ<sub>6</sub> model geomagnetic theodolite, yet they still await improvement.

In order for there to be a unified standard in China's station and field survey data, it was necessary to daily carry out absolute observation instrument constant determinations and comparisons using the Peking and Yushan Observatories as standards. At the same time, it was also necessary to improve the geomagnetic observation method and raise instrument precision, for example see references [42-46]. The use of the QHM nad BMZ instruments compare the instrument constants of each observatory and this has been carried out in the last few years as seen in reference [47a].

#### 8. Investigations and Theoretical Research of Seismo-Magnetic Relations

Because of the demands of earthquake prediction, there has been development of investigations and research of seismo-magnetic relations over the last ten years. A special feature was the development of experience in measuring magnetism and in prediction, such as the "magnetic storm two fold method"

"green and red light method" and the "magnetic storm nine fold method." However, there was the lack of a rational mechanism and model so that it was difficult to control and prove effective in practice. Attention was also given to industrial investigations and research. Reference [48] introduced the two analytical methods of "space interrelationship" and "phenomena interrelationship" for the use of geomagnetic field diurnal variation and based on the space interrelationship of the electromagnetic vertical weight of the Hongshan, Changli, Tianjin and Yushan Observatories and the Peking Observatory, the space interrelationship method was discussed for use in earthquake predictions. Reference [49] introduced the concept of the largest interrelated time and carried out preliminary research on the phase relations of the geomagnetic field vertical weight diurnal variations in two observatories. Related to the geomagnetic field diurnal variations, references [50-51], based on former work, introduced the local area geomagnetic diurnal variations analysis method and distinguished the geomagnetic local irregularities source field theory. At present, China also has several theories on the possible mechanism of earthquake relations. Reference [52] deduced a diffusion equation from Maxwell's equation which provided a basic equation of three dimensional electromagnetic induction, boundary conditions and a method for seeking solutions. Reference [53] considered that proceeding from the observed facts of the Xingtai earthquake in China and the geomagnetic variations, and because



they used the piezomagnetic effect to explain the seismo-magnetic relation, thus, like the earthquake "expansion" theory, there was introduced the idea of geomagnetic field variations induced by the minute cracking and expansion in rocks in an earthquake focus area. Yet, reference [53a] pointed out that this "expansion" effect is very small. Reference [54] used the magneto-elasticity theory and recommended the simultaneous following of the earthquake wave propagation whereby the magnetic field will produce a disturbance. Under ideal conditions in which the conductivity is  $\sigma \rightarrow \infty$ , the magnetic field variation in the earthquake focus region is about several gamma order of magnitude. This type of magnetic disturbance still exists in limited conductivity conditions. Reference [55] calculated the electromagnetic reaction of electric conducting spheres in electric conductance random variations and attempted to link "electromagnetic warnings" appearing before a strong earthquake. This is possibly one type of "warning mechanism." Other problems related to seismo-magnetic relations can also be seen in references [32-34]. Reference [56] summarized new developments related to structural magnetics introduced in the 1975 International Special Topics Meeting held in Paris.

## 9. Statistical Analyses of Solar-Terrestrial Relations

Early work on solar-terrestrial relations emphasized the statistical analyses of phenomena and here I will only give a

brief introduction. In recent years, the development of solar-terrestrial physics has been concentrated on research in the magnetosphere and aeronomy which are evaluated in another section of this article. References [58, 61-63] analyzed the relation of solar flares and magnetic storms. The relation of the position of solar flares on the sun's surface and magnetic disturbances was non-symmetrical and during the period 1938-1958 there were geomagnetic effects of large solar flares from the sun's  $\gamma$  and  $\beta\gamma$  sunspots in magnetic field areas. Reference [57] has statistics on the magnetic disturbance solar terrestrial magnetic level strength solar mean value from 1923-1954 and large magnetic storms averaged a 27 day period. Reference [59] used the corona green area 5303 Å materials from the United States' Climax Observatory during the 1909-1952 period and analyzed its relation with  $K_p$ . Reference [60] estimated the sun's average wind velocity as 300-500 kilometers/second for the relation of the flocculus and reappearance of magnetic interference during the periods 1942-1943 and 1952-1953. References [64-65] gave two examples for the relation of a solar eclipse and the upper atmosphere and stressed the influence of solar ultraviolet radiation changes in the ionosphere and ozone layer. Reference [66] used the 1955 edition of the Greenwich catalogue on large magnetic storms to analyze the gradual solar temperature changes at nine stations between 1909 and 1952 in China.

#### 10. Structure of Upper Atmosphere and Determination of Ozone Layer

China has still not systematically accumulated materials for determining the upper ozone layer. Up until recently, because the Doppler (?) spectrometer has been internationally corrected, two survey stations were established in Peking and in the southwest. References [67-70] discussed the reversal method and its improvement on the measurement of the vertical distribution of atmospheric ozone. There was great development in research on upper atmospheric structure, both in direct measurement and theoretical analysis. Direct measurement concentrated on less than 100 kilometers and used weather or sounding rockets to measure temperature, pressure and wind. For great altitudes, they used satellite orbit measurement and thus calculated upper atmospheric density and the earth's gravity [71-73]. In research on upper atmosphere types, references [74-76] discussed the dynamic transport and turbulent flow of the middle layer and heat layer atmosphere and their influence on the chemical composition following a high degree of distribution. References [77,77a] concerned with explanations and theories on the problems of upper atmosphere and dynamics, derived and discussed the heat atmosphere influence function (Green's function), the three layer layered type, the heat layer atmosphere tide differential operator and its eigenfunction and the diffusion wave in the upper atmosphere. References [78-82] analyzed and discussed two problems. For the first, there was analysis of the solar radiation heat

source on the heat layer atmosphere and they discussed the relation of diffusion action and upper atmosphere wave motion. They also established a diffusion equilibrium solution in an electromagnetic field and in a gravitational field and managed the distribution of a 500-50,000 kilometer ion composition. For the second, there was analysis of the radiation absorption and transmission process and heat equilibrium and they discussed the influence of early condensation, radiation, atmospheric escape and hothouse effect on planetary atmosphere and surface heat. It was considered that for the origin of the earth's and planetary atmosphere, the earth should have a very primitive atmospheric stage and that its main feature is its abundantly filled hydrogen and its compound reduced atmosphere. In the international geophysics year, China imported from abroad a whole set of night and daylight observing instruments, yet because of system and personnel changes, the addition of observation points has not been fixed and thus up until now it still has not been put into operation. On the theoretical analysis of airglow, reference [83] analyzed the half width of the oxygen green line altitude section and the material on the nearest atomic oxygen altitude section. The tendency was towards the excitation mechanism which did not resemble the three level reaction proposed by Professor S. Chapman but was a two pole reaction. Reference [84] overlooked radiation transition and absorption. They first calculated the electron excitation speed of the energy level in an oxygen atomic ground

state from the electron collision excitation cross-section and made hypotheses on the total collision excitation cross-section of neutral particles to attain a corresponding excitation speed. On this basis, they calculated the subenergy level excitation temperature in a 40-500 kilometer oxygen atom ground state.

#### 11. Ionospheric Measurement and Analysis

Because short wave radio communications are closely related to ionospheric conditions, early, serious attention was given in China to ionospheric measurement and analysis. It became even stronger after the international geophysics year (IGY). At present, domestic ionospheric frequency measurement, besides a minority of one or two stations, is basically man operated. Yet, measuring stations are numerous and their distribution is good and they have amassed long term source materials. The Chinese Electric Wave Propagation Institute designed and developed a completely automatic ionospheric altimeter and successively equipped each station. Moreover, they are in the process of speeding up work on frequency conversion in order to make a standard curve. For early research on ionospheric measurement see reference [85a]. In recent years, the Department of Physics in Wuhan University has manufactured an ionospheric automatic oblique return detector and the Space Physics Institute very recently designed, manufactured and researched an ionospheric electron capacity 505A model polarimeter and carried out receiving tests[85].

Early, domestic ionospheric analysis was also developed [86-89]. References [86] and [87] introduced the intersecting point correlative method and the ten point correlative method to analyze the statistical properties of vertically reflected quadratic echos for the ionosphere. References [88] and [89] analyzed the sunrise effect in the ionosphere's E and F layers and proved that the drift theory proposed by Dr. D.F. Madding was insufficient to explain the F<sub>2</sub> abnormality. During the 1977 and 1979 academic conferences, papers focused on aspects of the ionosphere showing the advances made in recent years. They can be divided into four areas: first was the sorting out of materials on vertical measurement and the conversion method of ionospheric electron concentration section structure [90-94]. Second was ionospheric frequency calculations, regular ionosphere and E<sub>s</sub> information channel properties and the problem of the ionospheric refraction effect [95-99]. Third was the analysis of features of certain Chinese area low ionosphere and ionospheric disturbances [100-104]. Fourth were references [105-111] which included the use of noise and other measures to calculate ionospheric parameters, the produced Doppler shift integral effect when high frequency electric waves pass ionospheric reflection, the influence of galactic cosmic rays on the ionosphere and the mutual action of electric waves, aerals and plasma.

## 12. Structure and Disturbance of Magnetosphere

Early work stressed theoretical analysis. References [112-114] touched upon the changes of external radiation band structures in magnetic storms and discussed a type of mechanism of charged particles passing into a geomagnetic field. References [115-116] focused on the qualitative appraisal of violent starting magnetic storms and the longitudinal non-symmetricality of the main phase of a magnetic storm. References [117-119] analyzed the movements of charged particles in geomagnetic fields and neutral line magnetic fields and they vigorously initiated simulated tests. References [120-121] summarized the propagation of magnetic fluid shock waves and based on the cosmic ray Forbush descent, they calculated the solar flare shock wave near the earth's orbit. China's putting a satellite into orbit created excellent conditions for space particle and magnetic field measurement and references [122-124] provide many examples representing this aspect of research. During the 1979 Solar-Terrestrial Physics Academic Conference, focus was placed on discussing solar eruptions, solar wind and magnetospheric substorms, especially the mechanism of magnetospheric substorms [125-127]. References [128-129] introduced a theoretical model of a satellite with a charged surface. References [130-131] discussed the geomagnetic field magnetic coordinates.

### 13. Studies on Cosmic Ray Intensity Variations

In the international geophysics year, China established observatories on Peking, Wuhan, Canton and Luoxue in Yunnan Province. However, at present, only the Peking Observatory maintains a neutron pile, an ionization chamber and a  $\mu$  meson telescope with three types of conventional observation. The other three observatories were soon discontinued. Early recording was good. For example, reference [132] wrote about the atmospheric pressure corrections of the cosmic ray neutron composition in the Peking and Luoxue Observatories. However, in recent years, for various reasons, regular observations and analyses still await reorganization and restoration. There have been a series of articles published on solar cosmic ray propagation. References [133-135] discussed the propagation of solar cosmic rays in interplanetary spaces including diffusion in interplanetary irregular magnetic fields and the two physical processes of solar wind and convection. For this, they used the non-dimensional constant to show particle diffusion and convection characteristics, sought a diffusion convection equation solution in a homogeneous infinite medium and attempted to use propagation correction in a solar proton event energy spectrum. References [136-138] took the interplanetary large scale fan shaped magnetic field as the basis to research the fan shaped coordinate system of solar cosmic rays in interplanetary propagation and derived



a propagation equation of solar cosmic rays in this type of coordinate system. Later, they further used the fan shaped coordinate system to discuss propagation and earth orbit modulation of Jupiter electrons in interplanetary space and set up a three dimensional diffusion model for solar cosmic ray diffusion. The reader is referred to reference [139] for the contents of cosmic ray intensity variations.

#### 14. Retrospect and Prospect

It can be seen from the above review that in the last thirty years China has made great advancements in geomagnetism and aeronomy and this has played a great role in building national defense and the national economy. On the one hand, the motive force for these advancements is owing to the task carrying the field of study and the field of study promoting the task. On the other hand, after determining long-range and annual plans for scientific and technological advancement, there has been continual support. However, many problems still exist, for example, the speed of development and the growth of personnel are too slow. In the last thirty years, there have only been 4-5 academic conferences related to this field (the first was in February, 1957, the second in September, 1963, the third in September, 1976, the fourth in February, 1979 and besides these there was a conference in November, 1977 which was limited to ionospheric physics). The proportion of articles in journals related to geomagnetism and

aeronomy is not very large and this cannot meet the needs of the "four modernizations." Furthermore, the development of each branch of geophysics has been unbalanced. During the 1950's, more stress was placed on geophysics, meteorology and seismology. After the international geophysics year and the beginning of the 1960's, there was development in geomagnetism and aeronomy. During the cultural revolution, all natural sciences, including geophysics, were effected by the impact of "ultraleftist" thought. In recent years, there has been notable development in geophysics, meteorology and especially seismology and when comparing, the development of geomagnetism and aeronomy depends on unified recognition and the adoption of measures for the impetuous to catch up.

At present, there is still a large gap between China and the advanced international level which is seen in basic construction, the use of advanced technology and the development of advanced original theories. Taking international magnetosphere study (IMS) as an example, in the vigorous development of international magnetosphere studies, the period 1976-1979 was one in which the whole world went from ground to outer space overall observation, and 1980-1981 is a period of data analysis and thorough research. When China first began work in this field, it depended on reorganizing China's various ground observations developed since the IGY and there are plans to develop the use

of new IMS equipment, carry out thorough research, make foreign things serve China, be based on the domestic and actively develop research on the magnetosphere and the solar-terrestrial relation [140]. Hereafter, in the area of basic construction, we should pay attention to maintaining the relative stability of the system of organization and industry, speed up the training of qualified personnel and quicken the filling in of the national deficiency in departments. It is also necessary to give attention to the quality of observatory materials and the fast processing of electronic computers. In the area of using advanced technology, we should pay attention to China's national conditions and economic effects. For example, in developing space technology and surveying, we can concentrate our strength on close areas, the small-scale or special characteristics and when study and achievement are joined we can at the same time tap the potential doors of the ground. In the area of theoretical research, we should promote the use of physics and mathematics theories to disclose the mechanism of geomagnetic and aeronomy phenomena and processes, utilize modern technology in understanding the basis of the microprocess to seek the macrolaw of natural change. At the same time, we should give attention to the mutual permeation of each discipline, expand and develop various types of discipline activities, strengthen domestic and international exchange and with continuous liberated thought vigorously cooperate and open up new roads in this branch of learning.

## Summary

# ON SOME ADVANCEMENT OF CHINESE GEOMAGNETISM AND AERONOMY DURING 1949—1979

Tschu Kang-kun

(Institute of Geophysics, Academia Sinica)

This paper gives a brief account of some advancement in the field of geomagnetism and aeronomy in China during the period of 1949—1979. The review covers the following 14 headings: (1) Establishment of geomagnetic observatories; (2) Geomagnetic surveys and charts; (3) Geomagnetic transient variation and geomagnetic storm predication; (4) Rock magnetism, paleomagnetism and archeomagnetism; (5) Magnetotelluric prospecting and related problems; (6) Magnetic prospecting and interpretation of data; (7) Some research on geomagnetic instruments; (8) Search of seismo-magnetic relations; (9) Statistical analyses of solar-terrestrial relations; (10) Structure of upper atmosphere and determination of ozone layer; (11) Ionospheric measurements and analyses; (12) Structure and disturbance of magnetosphere; (13) Studies on cosmic ray time variations; and (14) Retrospect and prospect.

## References

- [1] 余山地磁观测报告: 1957—1974; 北京地磁观测报告: 1957—1964; 广州地磁观测报告: 1956—1959; 兰州地磁观测报告: 1959—1961.
- [2] 陈宗器, 中国境内地磁观测之总检讨, 学术汇刊 1 (2), 99—120, 1944.
- [3] 中国科学院地球物理研究所, 地球基本磁场的分析方法, 1976.
- [4] 朱岗昆, 上海附近地磁场扰动变化的分析, (待发表).
- [5] 朱岗昆, 关于地球物理记录中之潮汐现象及其测定(英文), 地球物理学报, 2 (1), 74—82, 1950.
- [6] 朱岗昆等人, 北京地磁场和宇宙线强度的潮汐效应, 1963 年地球物理学会学术会议报告.
- [7] 朱岗昆, 关于太阳质子爆发的地球物理效应, 地球物理学报, 11(2), 183—208, 1962.
- [8] 高美庆, 郑体容, 地磁活动性预报工作初步总结, 1979 年日地物理学术会议报告.
- [8a] 赵学溥, 张树理, 146 MHz C 型爆发与地磁暴, 北京大学学报, 1978 年第 2 期.
- [8b] 侯民筠, 太阳爆发与地磁暴的多元相关分析, 北京大学学报, 1978 年第 3 期.
- [9] 许庆刚等 4 人, 中国某些地区岩石密度及磁化率资料的初步分析, 地球物理学报 9 (2), 149—154, 1966.
- [10] 侯重初, 大兴安岭地区的强磁性火山岩, 地球物理学报, 9 (2), 144—148, 1960.
- [11] 喻光楚, 介质的磁化率与包裹体形状的关系, 地球物理学报, 5 (1), 21—31, 1956.
- [12] 喻光楚, 介质的磁化率与包裹体浓度的关系, 地球物理学报, 6 (1), 69—76, 1957.
- [13] 张文佑, A. E. M. 斯恩, Some Palaeomagnetic Investigations on Chinese rocks, *Science Record*, 3 (1), 46—47, 1959.
- [14] 王子昌, 邓兴恩等, 应用中国岩石天然剩磁研究古地磁的初步结果, 地球物理学报, 9(2), 125—138, 1960.
- [15] 刘椿, 冯浩, 安徽休宁地区下震旦统休宁群砂岩的交变场退磁研究, 地球物理学报, 14 (3), 173—180, 1965.
- [16] 刘椿, 程国良等, 南京附近地区一些新生代玄武岩组的古地磁学研究及其地质年代的初步认识, 地球物理学报, 19 (2), 125—137, 1976.
- [17] 王普, 钱方等, 用古地磁方法对元谋人化石年代的初步研究, 中国科学, 第 6 期, 579—591, 1976.
- [18] 邓兴恩, 李东节, 北京地区史前地磁场及其变化的研究, 地球物理学报, 14 (3), 181—196, 1965.
- [19] K. 康达士, 1956 年 1 月 9—14 日在中国的北京和匈牙利的索波伦二处同时观测大地电流的结果, 地球物理学报, 5 (2), 79—92, 1956.
- [19a] K. 康达士, 地球物理勘探研究的新发展: 大地电流, 地球物理学报, 5 (2), 93—119, 1956.
- [20] A. F. 伊万诺夫, 中国石油普查地区的电法勘探, 地球物理学报, 8 (1), 71—83, 1959.
- [21] A. B. 卡文, 在中国几个地区进行的大地电流法勘探工作, 地球物理学报, 8 (2), 138—157, 1959.
- [22] 章冠人, 垂直旋转椭球体的自然电场及其解释, 地球物理学报, 5 (2), 177—182, 1956.
- [23] 章冠人, 水平旋转椭球体的自然电场及其解释, 地球物理学报, 5 (2), 183—188, 1956.
- [24] 章冠人, 倾斜旋转椭球体的自然电场及其解释, 地球物理学报, 6 (1), 87—94, 1957.

- [25] 章冠人, 当存在轴对称柱体时点电源的电场, 地球物理学报, 6 (2), 235—238, 1957.
- [26] 章冠人, 各向异性水平岩层对自然电场电位曲线的影响, 地球物理学报, 6 (1), 77—94, 1957.
- [27] 章冠人, 对三层电测深曲线等值原则证明的一些补充, 地球物理学报, 5 (1), 1—8, 1956.
- [28] 章冠人, 关于大地电流数据整理的总合法, 地球物理学报, 10 (2), 142—150, 1961.
- [28a] 雷海如, 对大地电流勘探法中扁圆法的改进——兼论大地电流观测资料整理的新方向, 1963年地球物理学会学术会议报告.
- [28b] 雷海如, 关于整理大地电流观测资料的总和法, 地球物理学报, 13 (3), 227—233, 1964.
- [29] 李佩霞, 大地电流法中垂直断层的正演问题, 地球物理学报, 10 (1), 79—82, 1961.
- [30] 李佩霞, 大地电流法中倾斜断层的正演问题, 地球物理学报, 11 (2), 178—182, 1962.
- [31] 张河墨, 利用大地电流法研究垂直断层, 地球物理学报, 11 (2), 170—177, 1962.
- [32] 兰州地震大队大地电磁深探组, 中国南北地震带北段内地壳和上地幔的电性特征, 地球物理学报, 19 (1), 28—34, 1976.
- [33] 陈伯豹, 渤海西岸的电导率异常, 地球物理学报, 17 (3), 169—172, 1974.
- [34] 余文燮, 祁锡、王仕明, 甘肃东部地区短周期地磁变化异常及其与地震的关系, 地球物理学报, 21 (3), 218—224, 1978.
- [35] G. Portstendort, “大地电磁勘探原理” 1975, (中国科学院地球物理所中译本, 1976年8月).
- [36] 熊光楚, 金属矿区地面磁法研究工作中的几个问题, 地球物理学报, 15 (1), 1—5, 1966.
- [36a] 熊光楚等, 近似积分在磁法数据推断中的应用, 1963年地球物理学会学术会议报告.
- [36b] 熊光楚等, 用磁法法求标本时几个计算标本磁化强度的公式, 1963年地球物理学会学术会议报告.
- [37] 余世新, 二度磁场解析方法, 地球物理学报, 14 (2), 136—146, 1965.
- [38] 熊光楚等, 某铁矿磁异常的定量推断, 地球物理学报, 15 (1), 6—13, 1966.
- [39] 黄树棠, 复杂磁异常的定量解释方法, 地球物理学报, 15 (1), 33—52, 1966.
- [40] 侯重初, 关于  $\Delta T$  磁异常的解算, 地球物理学报, 13 (4), 324—327, 1964.
- [40a] 侯重初, 斜磁化板状体产状单元的计算方法, 1963年地球物理学会学术会议报告.
- [40b] 侯重初, 确定水平圆柱体磁化倾角的简易方法, 1963年地球物理学会学术会议报告.
- [41] 段星北, 李平一, 一种地磁( $\Delta Z$ )化极方法, 地球物理学报, 22 (2), 169—180, 1979.
- [42] 陈志强, 关于水平强度H磁变仪的装置(英文), 地球物理学报, 2 (1), 83—86, 1950.
- [42a] 胡岳仁, 磁通量平面的一个方法(英文), 地球物理学报, 2 (3), 299—303, 1951.
- [43] 刘庆令, 黄锡令, 徐元芳, 磁致温度系数及感应系数的测定, 地球物理学报, 10 (2), 126—131, 1961.
- [44] 顾子明, 詹雨律, 地磁目视记录仪准确度的提高, 地球物理学报, 13 (1), 24—28, 1964.
- [45] 顾子明, 黄锡令, 磁通门式磁仪测磁倾角的方位误差, 地球物理学报, 22 (1), 84—88, 1979.
- [46] 地球物理所电法勘探组, 大地电流勘探用的调制式直流放大器, 地球物理学报, 8 (2), 175—184, 1959.
- [47] 孙仿友等, 我国各地磁台 QHM 比测, 1963年地球物理学会学术会议报告.
- [47a] 孙仿友等, 地磁台仪器比测工作总结, 1975年.
- [48] 中国科学院地球所 10室1组, 地磁场的空间相关性及其在地磁预报中的应用, 地球物理学报, 20 (3), 169—184, 1977.
- [49] 陈冠晃, 王鼎盛, 两台地磁场垂直分量日变形态的位相关系分析, 地球物理学报, 22 (1), 56—65, 1979.
- [50] 祁贵仲, 局部地区地磁日变分析方法及中国地区  $S_q$  场的经度效应, 地球物理学报, 18 (2), 104—117, 1975.
- [51] 祁贵仲, 区分地磁局部异常的源场理论, 地球物理学报, 22 (1), 66—77, 1979.
- [52] 中国科学院地球所 10室2组, 地震的感应磁效应, 地球物理学报, 20 (1), 70—80, 1977.
- [53] 祁贵仲, “膨胀”磁效应, 地球物理学报, 21 (1), 18—33, 1978.
- [53a] 徐世新, 关于压磁效应和膨胀效应, 地震学报, 1 (1), 76—81, 1979.
- [54] 王水, 郭自强, 林启朝, 伴随地震波动的一种磁扰, 地球物理学报, 22 (1), 78—83, 1979.
- [55] 李钧, 导电率随时间变化的导电球的电磁响应, 地球物理学报, 22 (3), 243—254, 1979.
- [56] 朱尚昆, 关于大地构造磁学和构造电学的新进展, 国外地质, 第七期, 17—22, 1977.
- [57] 胡岳仁, 太阳自转对地球磁场的影响, 地球物理学报, 7 (2), 103—107, 1958; 8 (2), 123—131, 1959.
- [58] 蒋伯琴, 太阳耀斑与磁暴的关系, 地球物理学报, 3 (1), 38—46, 1960.
- [58a] 汪瑜容等, 地球磁暴与太阳耀斑的关系, 1963年地球物理学会学术会议报告.
- [59] 刘传新, 日冕绿区与地磁扰动间的某些问题, 地球物理学报, 10 (1), 17—26, 1961.
- [60] 郭亨, 关于1942—1943, 1952—1953年谱斑与重现磁扰的关系, 地球物理学报, 10 (2), 151—159, 1961.
- [61] 章公亮, 太阳活动对地磁场影响的南北不对称性, 地球物理学报, 11 (1), 92—94, 1962.
- [62] 章公亮, 耀斑日面位置与磁扰关系的不对称性, 地球物理学报, 12 (1), 32—40, 1963.
- [62a] 方成, 蒋晓霞, 耀斑日面位置和磁扰的关系, 1963年地球物理学会学术会议报告.
- [62b] 章公亮, 刘其俊, 磁扰 Kp 类型与耀斑活动(1935—1958), 1979年地物理学会会议报告.
- [63] 刘传新, 太阳  $\tau$  及  $\beta\gamma$  黑子磁区大耀斑的地磁效应, 地球物理学报, 13 (1), 11—19, 1961.
- [64] 莫纪华, 日食与电离层的关系, 地球物理学报, 3 (1), 9—21, 1954.
- [64a] 赵崇路, 1968年日全食电离层观测资料分析, 1977年电离层会议报告.

- [65] H. H. 巴里斯基, 胡仁超等 4 人, 1958 年 4 月 19 日海南岛日环食臭氧层的变化, 地球物理学报, 10 (1), 1—16, 1961.
- [66] 杨鉴初, 大磁暴后我国磁度的变化, 地球物理学报, 10 (2), 101—112, 1961.
- [67] 魏鼎文, Gutz 逆转效应中 13112 Å 的有效散射高度上升过程的跳跃性及逆转方法 B 解的不唯一性, 地球物理学报, 11 (2), 123—135, 1962.
- [68] 李利斌, 黄信楠, 李钧, 关于逆转方法 B 解的不唯一性, 地球物理学报, 13 (1), 20—23, 1964.
- [69] 魏鼎文, 林启锦, 测量大气臭氧垂直分布的逆转方法 (C) 对逆转方法 (A)、(B) 的改进, 地球物理学报, 13 (4) 260—272, 1964.
- [70] 魏鼎文, 再谈关于逆转方法 (B) 解的不唯一性, 地球物理学报, 13 (4), 329—332, 1964.
- [71] 朱岗昆等, 人造卫星的运动及其在地球物理学中的应用, 1963 年天文学应用会议报告.
- [72] 中国科学院人卫观测系统, 290 公里高度附近大气密度及其变化的测定, 1979 年日地物理学学术会议报告.
- [73] 刘振兴, 流星和高层大气分子间相互作用的热力学和动力学过程, 地球物理学报 11 (1), 1—11, 1962.
- [74] 陈耀武, 对高层大气模式的研究, 地球物理学报, 12 (1), 1—11, 1963.
- [75] 陈耀武, 中层和热层大气的动力传输及湍流系数随高度的分布, 1979 年日地物理学学术会议报告.
- [75a] 刘振兴, 中层和热层大气的湍流结构, 1979 年日地物理学学术会议报告.
- [76] 陈耀武, 中层和热层大气的动力传输及其对化学成分随高度分布的影响, 1979 年日地物理学学术会议报告.
- [77] 陈哲明, 高层大气结构与动力学若干问题的解释理论, 中国科学, 第 4 期, 416—429, 1978.
- [77a] 陈哲明, 热层大气潮汐微分算子与微分方程, 地球物理学报, 22 (4), 1979.
- [78] 陈哲明, 热层大气中的扩散波, 1979 年日地物理学学术会议报告.
- [79] 陈哲明, 太阳辐射等热流对热层大气的影响, 1979 年日地物理学学术会议报告.
- [80] 陈哲明, 马式明, 等离子体层下部 500—50000 公里电离成分的分布, 1979 年日地物理学学术会议报告.
- [81] 陈哲明, 马式明, 行星高层大气的辐射吸收传递过程和热平衡, 1979 年日地物理学学术会议报告.
- [82] 陈哲明, 地球与行星大气的起源, 天文学报, 19 (1), 81—91, 1978.
- [83] 傅鸿生, 高层大气中氧绿线强度与氧原子密度的关系, 地球物理学报, (待发表).
- [84] 吴健征, 热层氧原子基态中  $^3P_1$  亚能级的激发温度, 1979 年日地物理学学术会议报告.
- [85] 陈培人, 丘雪明, 葛和荣, 徐楚孚, 利用同步卫星信标研究电离层, 地球物理学报, (待发表).
- [85a] 周纯, 电离层的直接探测研究, 地球物理学报, 12 (2), 100—117, 1963.
- [86] 李钧, 相关分析法在研究电离层不均匀结构中的应用, 地球物理学报, 11 (2), 105—122, 1962.
- [87] 李钧, 从电离层垂直反射的二次回波产生特征, 地球物理学报, 14 (2), 85—93, 1965.
- [88] 张肇先, 电离层 E 层和 F 层中的日出效应, 地球物理学报, 12 (1), 12—31, 1963.
- [89] 李维祺, 迁移理论解释 F<sub>2</sub> 异常的可能性, 地球物理学报, 15 (2), 129—141, 1966.
- [89a] 肖佐, 傅竹风, 流磁波在电离层中的传播, 1963 年地球物理学学术会议报告.
- [90] 何友文, 联合运用二类特征波计算电离层电子浓度垂直分布的一种方法, 1977 年电离层会议报告.
- [91] 吴均英, 电离层高频记录分析方法, 1977 年电离层会议报告.
- [92] 张肇先, 夜间电离层高频曲线的一种新的换算方法, 地球物理学报, (待发表).
- [93] 刘培静, F<sub>2</sub> 层临界频率  $f^oF_2$  月中值统计, 1977 年电离层会议报告.
- [94] 刘培静, 宋英亭, 电离层电子浓度平均剖面的提供方法, 1977 年电离层会议报告.
- [95] 孙克儒, F<sub>2</sub> 层局部地区频率预测方法, 1977 年电离层会议报告.
- [96] 焦培南, 短波电离层无线电通讯系统性能的预测, 1977 年电离层会议报告.
- [97] 张训斌, 电离层短波信道特性测量, 1977 年电离层会议报告.
- [98] 王椿年, 电离层模式预报的一种方法, 1977 年电离层会议报告.
- [99] 张仁芳, 电离层折射指数级数表示法的估算, 1977 年电离层会议报告.
- [100] 李赞红, 武昌上空电离层夜间 E 层, 1977 年电离层会议报告.
- [101] 叶永寿, 武昌地区电离层暴的初步研究, 1963 年地球物理学学术会议报告.
- [102] 叶永寿, 在地磁暴和宇宙线暴期间 D 层的扰动及其对无线电波传播的影响, 1979 年日地物理学学术会议报告.
- [103] 北大  $\times \times \times$ , 关于中纬电离层暴开始时间及形态的分析, 1976 年日地物理学学术会议报告.
- [104] 索玉成, 宋英亭等, 海南、荆州地区电离层暴扰的某些特征, 1979 年日地物理学学术会议报告.
- [105] 郑荣华, 用哨声推算电离层的电子密度, 1977 年电离层会议报告.
- [105a] 何友文, 我国低电离层的观测与分析, 1979 年日地物理学学术会议报告.
- [106] 王水, 计算电离层 F 边界反射系数的一种方法, 1977 年电离层会议报告.
- [107] 郑传青, 利用最小延迟曲线计算电离层参数, 1977 年电离层会议报告.
- [108] 黄雪钦, 电离层非相干散射探测中的信号积累问题, 1977 年电离层会议报告.
- [109] 龙成灵, 侯杰昌, 关于电离层反射电波频率变化的问题, 地球物理学报, 22 (4), 1979.
- [110] 夏培瑞, 稳定均匀的等离子体板对电波传播和反射特性的影响, 1979 年日地物理学学术会议报告.
- [110a] 王文清, 肖佐, 雷宏通, 葛相仁, 等离子体鞘套对天线阻抗及辐射性能的影响, 1976 年日地物理学学术会议报告.
- [110b] 肖佐, 余辉等离子体中的谐振探针实验, 地球物理学报, (待发表).
- [111] 王作胜, 关于银河宇宙线在电离层 D 层中电子产生率的计算, 1979 年日地物理学学术会议报告.

- [112] 濮祖荫、李文艺, 磁暴主相期外辐射带结构的变化, 地球物理学报, 13 (2), 91—105, 1964.
- [112a] 刘振兴、濮祖荫, 磁暴期间外辐射带结构的变化, 地球物理学报, 13 (3), 189—200, 1964.
- [113] 徐荣兰、赵九章, 磁暴扰动期间电离层捕获区的变化, 地球物理学报, 11 (1), 12—21, 1962.
- [114] 都亨、赵九章, 带电粒子穿入地磁场的一种机制, 地球物理学报, 13 (3), 201—210, 1964.
- [115] 刘传群、谷小翠, 急始型磁暴的定性估计, 地球物理学报, 13 (4), 249—259, 1964.
- [116] 都亨, 磁暴主相的经度不对称性, 地球物理学报, 13 (3), 211—215, 1964.
- [117] 徐荣兰、周国成, 带电粒子在磁场中运动区及其模型实验, 地球物理学报, 12 (2), 121—130, 1963.
- [118] 徐荣兰、刘梅香, 地磁捕获区及某些空间相关现象的非轴对称形状, 地球物理学报, 14 (2), 73—84, 1965.
- [119] 徐荣兰、范振福, 带电粒子在中性线磁场中的运动, 1979 年日地物理学术会议报告.
- [120] 章公亮, 磁流体冲击波在非均匀磁场中的传播, 地球物理学报, 13 (1), 1—10, 1964.
- [121] 魏春思, 根据宇宙线 Forbush 下降计算地球轨道附近的耀斑击波, 地球物理学报(待发表).
- [122] 王作柱、黄永年、周国成等, 一种空间半导体质子探测器, 能谱计算和校准, 半全同探测器计数率和投掷角分布, 1979 年日地物理学术会议报告.
- [122a] 韦瑞干等, 卫星闪烁计数器几个参数的确定, 1979 年日地物理学术会议报告.
- [123] 赖世道、王惠民, 卫星磁偶极矩的测量, 1979 年日地物理学术会议报告.
- [124] “人造地球卫星环境手册”, 国防出版社, 1971.
- [125] 胡文瑞, 太阳风湍流和磁层亚暴机制, 1979 年日地物理学术会议报告.
- [126] 刘振兴, 磁层亚暴期间的离子声流异常电阻和粒子的加速, 1979 年日地物理学术会议报告.
- [127] 刘振兴, 磁层中性线区的宏观动力学过程与磁层亚暴, 1979 年日地物理学术会议报告.
- [128] 刘振兴等, 磁层亚暴期间同步高度电子通量突增的理论计算, 1979 年日地物理学术会议报告.
- [129] 丘雪明, 同步轨道卫星表面带电的理论模型, 1979 年日地物理学术会议报告.
- [130] 刘振兴, 磁坐标的简化计算及其应用, 1979 年日地物理学术会议报告.
- [131] 周国成, 源壳壳分离和磁坐标, 1979 年日地物理学术会议报告.
- [132] 寿孟岗等人, 北京、乌鲁木齐地宇宙线强度气压系数的初步分析, 1963 年地球物理学会学术会议报告.
- [133] 章公亮, 用量纲分析法求解太阳宇宙线扩散对流方程, 高能物理与核物理 2 (3), 200—209, 1978.
- [134] 章公亮, 太阳宇宙线的传播效应, 高能物理与核物理, 2 (5), 285—293, 1978.
- [135] 章公亮, 太阳质子事件能谱的传播改正, 1979 年日地物理学术会议报告.
- [136] 黄永年, 关于太阳宇宙线传播问题的研究, 地球物理学报 22 (2), 97—108, 1979.
- [137] 黄永年, 关于木星电子在行星际空间的传播及地球轨道上的调制, (待发表).
- [138] 黄永年, 太阳宇宙线传播的三维扩散模式, 1979 年日地物理学术会议报告.
- [139] Л. И. Дорман, “宇宙线强度变化”, 科学出版社, 1965 年.
- [140] 朱岗昆, 关于国际磁层研究 (IMS) 和我们的初步打算, 1979 年日地物理学术会议报告.

END

DATE  
FILMED

5 81

DTIC